

Shot or Sprouted Grain Classification Procedure

Background

Rain immediately before or during harvest can affect grain quality and have a negative impact on the end-use product. For example, shot wheat can lead to flour failing to rise during baking, while sprouted barley may fail to germinate during the malting process.

A Falling Number machine is used to measure the impact of shot wheat or sprouted barley on the quality of the grain. High quality grain will have a higher falling number.

Barley may also be tested using a Rapid Viscosity Analyser (RVA) with malting barley requiring a result of more than 130 units.

The Falling Number and RVA results for different wheat and barley grades are:

Commodity	Grade	Falling Number (seconds)	RVA (units)
Wheat	H1, H2, APW, ASW, DR1, DR2	300 or above	NA
	AUW, AUH	250 or above	
	AGP, DR3	200 or above	
	Feed	No Limit	
Barley	Malting	300 or above	130 or above
	Feed	No Limit	No Limit

Shot or Sprouted Grain Classification Procedure

- Wheat and malting barley samples with sprouted or shot grains will be assessed using a Falling Number machine. A Rapid Viscosity Analyser (RVA) may be used for assessing malting barley.
- A retest may be requested, as per the [Classification Dispute Resolution Procedure](#), where the first test has the following results:

Commodity	Falling Number or RVA test	Retest option
Wheat	Falling Number	Within 15 seconds
Barley	Falling Number	Within 15 seconds
Barley	RVA	Within 10 units

- Grain samples are routinely tested post-delivery; with the results possibly identifying the need to implement Falling Number or RVA tests.
- Growers may forgo a Falling Number or RVA test, accepting a feed grade.